

**ROACH & ANT GEL SYRINGES**

*Bait Gel*

**ROACH GEL SYRINGES** are the most effective and fastest roach bait available. Consumption is not actually necessary, cockroaches only have to touch the bait to die.

**ANT GEL SYRINGES** provide fast control of ants within three to five days. Its active ingredient, Fipronil, allows foraging ants to return to the colony and feed the bait to larvae and the queen(s) so that the entire colony is killed.

**DIRECTIONS FOR USE:**

**ROACH GEL SYRINGES:** Application rate is dependent on level of infestation and species to be controlled, see

application rate table on label. Apply bait as a spot or as a bead in cracks and crevices. The lower application rate should be used for low to moderate infestations or re-treatments, and the higher rate for more severe infestations or where the population occurs in

should be at or near as corners, or cracks and inaccessible areas. Gel placements harborages or aggregation areas, such crevices. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control. Place bait under baseboards, sinks, countertops, seats, around water pipes, water heaters, dishwasher equipment, in cracks and crevices as well as voids where insects harbor. For outdoor use (adjacent to home -or- structure): Place bait in areas adjacent to structures where roaches may nest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints. Apply bait in protected areas whenever possible. Environmental stresses such as direct sunlight and water will reduce the residual effectiveness. Priority treatment areas may be identified by visual inspection of such areas listed above or aided by trapping techniques or spot flushing.

**ANT GEL SYRINGES:** Application rate is dependent on level of infestation and species to be controlled. Bait should be applied as spots. Applications in Food/Feed Handling Establishments. Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only. Apply with syringe applicator or bait injector directly into cracks and crevices. Place applicator tip into cracks and crevices and where ants may travel. Apply gel (up to 1 gram) per spot. Apply bait in small amounts between different elements of construction, openings



**ROACH GEL SYRINGES**  
**ACTIVE INGREDIENT:**  
 Fipronil\*: [5-Amino-1-(2, 6-dichloro-4-trifluoromethyl phenyl)-4-(1, R, S)-(trifluoromethyl sulfinyl)-1H-pyrazole-3-carbonitrile]:.....0.01%  
**OTHER INGREDIENTS:**.....99.99%  
 Total:.....100.00%  
 \*CAS NO. 120068-37-3  
**PRODUCT # 9286**

**ANT GEL SYRINGES**  
**ACTIVE INGREDIENT:** Fipronil\*:.....0.001%  
**OTHER INGREDIENTS:**.....99.999%  
 Total:.....100.000%  
 CAS NO. 120068-37-3  
**PRODUCT # 7441**

**FOR PROFESSIONAL USE ONLY**  
 CONSULT LABEL AND SDS BEFORE USING.



leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, cabinets, sinks, stoves and other places where ants may enter the establishment. During follow-up visits inspect bait placements and reapply when necessary. Do not apply bait to areas where food/feed, food utensils or food processing surfaces may become contaminated. Do not apply bait in cracks and crevices of areas that are routinely washed, tops of tables, food/feed preparation and prepared food/feed holding surfaces as bait may be removed by washing. Do not apply bait to surfaces where the temperature exceeds 120° F as bait may run. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove it and wash exposed surface.

**ROACH & ANT GEL SYRINGES**